

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12172 A1

(51) International Patent Classification⁷: A61K 31/00,
39/395, 49/00, C07K 2/00, 5/00, 16/28

(21) International Application Number: PCT/US00/22670

(22) International Filing Date: 17 August 2000 (17.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/149,258 17 August 1999 (17.08.1999) US

(71) Applicant: PURDUE RESEARCH FOUNDATION
[US/US]; 1291 Cumberland Avenue, West Lafayette, IN
47906 (US).

(72) Inventor: KINCH, Michael, Scott; 2320 Flowering Crab
Drive, Lafayette, IN 47905 (US).

(74) Agent: LAMMERT, Steven, R.; Barnes & Thornburg, 11
South Meridian Street, Indianapolis, IN 46204 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,

DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.
- With (an) indication(s) in relation to deposited biological
material furnished under Rule 13bis separately from the
description.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 01/12172 A1

(54) Title: TREATMENT OF METASTATIC DISEASE

(57) Abstract: The present invention is directed to compounds and methods for the treatment of metastatic disease. The compounds of this invention have specificity for EphA2, an epithelial cell tyrosine kinase that is overexpressed in metastatic tumor cells. The compounds used in accordance with this invention may be provided in a pharmaceutical composition for treatment of metastatic disease.

BEST AVAILABLE COPY

TREATMENT OF METASTATIC DISEASE

Field of the Invention

The present invention relates to diagnosis and treatment of metastatic disease. More particularly, this invention relates to the use of an extracellular epitope of an epithelial cell tyrosine kinase that is overexpressed in metastatic tumor cells as the target for the diagnosis and treatment of metastatic disease. Most particularly, this invention relates to the use of compounds that interact with and alter expression of the epithelial cell tyrosine kinase.

10

Background and Summary of the Invention

Cancer is a disease of aberrant signal transduction. The most dangerous forms of cancer are malignant cells that metastasize to distant sites in a body. Metastatic cells have acquired the ability to break away from the primary tumor, translocate to distant sites, and colonize distant and foreign microenvironments. Cancer cell metastasis requires cellular capacity to 1) detach from a primary tumor, 2) migrate and invade through local tissues, 3) translocate to distant sites in the body (via lymph or blood), 4) colonize a foreign site, and 5) grow and survive in this foreign environment. All of these behaviors are linked to cell adhesions. Cell adhesions control the physical interactions of cells with their microenvironment. Cell adhesions also initiate signals that dictate tumor cell growth, death, and differentiation. At the cellular level, metastatic cells have overcome restraints upon cell growth and migration that result from physical linkages and signals conveyed by cell-cell contacts. Malignant cells often have increased interactions with surrounding extracellular matrix (ECM) proteins that provide linkages and signals that promote several aspects of metastasis.

Levels of protein tyrosine phosphorylation regulate a balance between cell-cell and cell-ECM adhesions in epithelial cells. Elevated tyrosine kinase activity weakens cell-cell contacts and promotes ECM adhesions. Alteration in levels of tyrosine phosphorylation is believed to be important for tumor cell invasiveness. Tyrosine phosphorylation is controlled by cell membrane tyrosine kinases, and increased expression of tyrosine kinases is known to occur in metastatic cancer cells.

EphA2 is a 130 kDa receptor tyrosine kinase that is expressed on adult epithelia. A member of the Eph family of tyrosine kinases known as Ephrins, EphA2 is a transmembrane receptor tyrosine kinase with a cell-bound ligand. EphA2 expression has been found to be altered in many metastatic cells, including lung, breast, colon, and prostate tumors. Additionally, the distribution and/or phosphorylation of EphA2 is altered in metastatic cells. Moreover, cells that have been transformed to overexpress EphA2 demonstrate malignant growth, and stimulation of EphA2 is sufficient to reverse malignant growth and invasiveness. EphA2 is a powerful oncoprotein. The present invention is directed to compounds and methods that target EphA2 for the diagnosis and treatment of metastatic cancers.

One approach to cancer therapy is the administration of preformed antibodies to predetermined tumor antigens. This process is known as passive antibody treatment. An example of passive antibody treatment is the use of Herceptin® for the treatment of breast cancer. Herceptin® is a humanized form of a murine monoclonal antibody specific for the extracellular domain of Her2/Neu. The basis for treatment with Herceptin® is that 25-30% of metastatic breast cancers overexpress the Her2/Neu receptor tyrosine kinase. Herceptin® has been well tolerated in clinical trials and shows much promise for the maintenance and regression of metastatic breast cancer.

Effective passive immunotherapy for treatment of tumors requires isolation and preparation of an antibody that: 1) targets an antigen that is overexpressed in metastatic tumors; 2) targets an extracellular epitope of said antigen; 3) is not cross-reactive with any other antigen in a patient's circulation; and 4) exhibits tumoricidal or tumoristatic activity.

In a preferred embodiment, this invention relates to the selection and use of antibodies that are specific to an extracellular epitope of EphA2. The methods of this invention include the preparation, selection, and use of EphA2 specific antibodies for cancer therapy.

Another approach to cancer treatment is to use agonists to stimulate expression. For example, EphrinA1-F_c, the extracellular domain of ephrinA1 linked to immunoglobulin heavy chain (see Miao, H., et al., EphA2 kinase associates with focal adhesion kinase and upon activation, inhibits integrin-mediated cell adhesion

and migration, Nature Cell Biol 2, 62-69 (2000)), can be used to increase the phosphotyrosine content of EphA2. Thus, in another preferred embodiment, this invention relates to the use of agonists or antagonists to alter the expression of EphA2 in metastatic cells.

5 Thus, this invention is directed to the use of agonists and antagonists to alter the expression of EphA2. EphA2 may be targeted by use of artificial or hybrid forms of the protein, protein inhibitors, antisense oligonucleotides, or small molecule inhibitors. Also, while a preferred embodiment is directed to use of monoclonal antibodies, polyclonal, artificial, and hybrid antibodies are known in the art. It should
10 be understood that use of techniques known in the art to target EphA2 are within the scope of this invention.

 One aspect of this invention is a pharmaceutical composition for the treatment of mammalian metastatic tumors which overexpress EphA2, comprising a compound that specifically interacts with an extracellular epitope of EphA2 in an
15 amount effective to reduce metastatic disease and a pharmaceutically acceptable carrier. In the preferred embodiment, the pharmaceutical composition comprises the antibody B2D6, an antibody that specifically binds to an extracellular epitope of EphA2

 Another aspect of this invention is a method of treating a patient
20 having a metastatic tumor which overexpresses EphA2. The method comprises administering to the patient a therapeutic amount of a compound that binds to an extracellular epitope of EphA2. In a preferred embodiment, the compound is an antibody.

 A third aspect of this invention is a method for detecting the presence
25 of metastatic cells. The method includes use of a labeled antibody specific to an extracellular epitope of EphA2. A cell sample is incubated with the antibody, unbound antibody is removed, and the bound labeled antibody is detected.

 An additional aspect of this invention is a method for producing antibodies which inhibit metastatic tumor proliferation by specifically binding to an
30 extracellular epitope of EphA2. This method includes injecting tyrosine phosphorylated proteins into the lymph nodes of a mammal, harvesting the lymph

nodes, fusing the lymph node cells with myeloma cells to form hybridomas, and selecting hybridomas which produce antibodies specific for EphA2.

In still another aspect of this invention, a pharmaceutical composition for treatment of a mammalian metastatic tumor is provided, the composition
5 comprising a compound that alters expression of EphA2 in an amount effective to reduce metastatic proliferation of said tumor, and a pharmaceutically acceptable carrier therefor. In a preferred embodiment, the composition comprises an ephrin.

Additional features of the present invention will become apparent to those skilled in the art upon consideration of the following detailed description of
10 preferred embodiments exemplifying the best mode of carrying out the invention as presently perceived.

Brief Description of the Drawings

Fig. 1 is an overview of the RIMMS procedure, through which the
15 antibodies of this invention are generated;

Fig. 2A-C show a series of western blots showing EphA2 expression in human cell lines;

Fig. 2A is a western blot showing EphA2 expression in various human prostate cancer cell lines;

20 Fig. 2B is similar to Fig. 2A, except showing EphA2 expression in a human prostatic epithelial cell line and expression in the cell line after transformation by oncogenic K-Ras or X-irradiation;

Fig. 2C is similar to Fig. 2B, except showing EphA2 expression in another human prostatic epithelial cell line and expression in the cell line after
25 transformation by oncogenic K-Ras or X-irradiation;

Fig. 3 is similar to Fig. 2, except showing EphA2 expression in canine prostatic cancer cells; and

Fig. 4 shows predicted antibody binding plotted against cell density in a screening procedure for antibodies which are specific for an extracellular epitope of
30 EphA2.

Detailed Description of the Invention

EphA2 is expressed differently in normal and metastatic cells. In normal breast and prostate epithelial cells, EphA2 is enriched in within cites of cell adhesion. Conversely, in metastatic prostate cells EphA2 is diffusely distributed, and
5 in metastatic breast cancer cells EphA2 is redistributed into the membrane ruffles. EphA2 expression is also known to be altered in lung and colon malignancies, and it is believed that EphA2 altered expression occurs in other types of metastasis, particularly epithelial malignancies. Thus, techniques designed to alter EphA2 expression can be exploited to diagnose and treat metastatic disease.

10 In a preferred embodiment, antibodies specific for tyrosine phosphorylated proteins in cancer cells have been isolated and used to target cancer cells in passive immunotherapy. This approach is based upon the fact that many tyrosine kinases, e.g., Her2/Neu, are expressed by oncogenes and are therefore overexpressed in cancer cells. The present invention is directed to the production and
15 use of antibodies capable of recognition of and specific binding to extracellular epitopes of the tyrosine kinase EphA2. The antibodies are produced by selected hybridomas, themselves the product of fusion of myeloma cells with lymph node cells harvested from animals subjected to a specific inoculation protocol designed for increased sensitivity and diversity of responding hybridomas.

20 To produce these hybridomas, tyrosine phosphorylated proteins from Ras-transformed human epithelial cells were isolated by affinity chromatography using existing phosphotyrosine specific antibodies. The tyrosine phosphorylated proteins are then used as an immunogen for producing monoclonal antibodies according to the procedure illustrated in Fig. 1. Low-dose amounts of tyrosine
25 phosphorylated proteins are injected proximal to lymph nodes of a mammal, every other day, over a ten day period (the RIMMS strategy). B cells from engorged lymph nodes are then isolated and fused with Bcl-2-overexpressing myeloma cells, to minimize apoptosis after fusion. This method results in increased diversity, specificity, and cost-effectiveness of hybridoma production. The hybridomas are
30 screened to identify those hybridomas producing antibodies that distinguish malignant from normal cancer cells.

Hybridomas producing antibodies specific to EphA2 have been selected. Use of the RIMMS technique has resulted in the production of a multiplicity of hybridomas producing monoclonal antibodies that specifically bind EphA2. To date, at least 450 hybridomas have been identified which produce antibodies capable of distinguishing malignant from normal cancer cells. Of the first four such hybridomas to be characterized, two recognize independent epitopes on EphA2. The first, D7, produces an antibody recognizing an intracellular epitope. The second, B2D6, produces an antibody that specifically binds an extracellular epitope of EphA2, a characteristic that enables its effective use for the diagnosis and treatment of selected metastatic tumors.

While the RIMMS strategy has proven to be valuable in the production of EphA2 specific antibodies, other techniques are known in the art for producing antibodies to a specific antigen, and these techniques are within the scope of this invention.

It is known in the art to use antibodies to detect the presence or overexpression of a specific protein. Because EphA2 is overexpressed in metastatic cells, EphA2-specific antibodies of this invention may be used to detect this overexpression and, thus, to detect metastatic disease. Such techniques include but are not limited to western blotting, precipitation, agglutination, and ELISA assays. These techniques are well known in the art. Also, the extracellular epitope specificity of EphA2-specific antibodies of this invention can be exploited to detect changes in EphA2 localization which are associated with metastasis. In normal breast and prostate epithelial cells, EphA2 is enriched in within cites of cell adhesion, whereas in metastatic cells, EphA2 distribution is altered. In metastatic prostate cells EphA2 is diffusely distributed, and in metastatic breast cancer cells EphA2 is redistributed into the membrane ruffles. EphA2 expression is also known to be altered in lung and colon malignancies, and it is believed that EphA2 altered expression occurs in other types of metastasis, particularly epithelial malignancies. Techniques such as immunohistological staining or immunofluorescent microscopy are well known in the art and may be used to visualize EphA2 distribution. See, for example, U.S. Patent No. 5,514,554. In order to detect overexpression or altered distribution of EphA2, the EphA2-specific antibodies may be labeled covalently or non-covalently with any of a

number of known detectable labels, such fluorescent or radioactive substances, as is known in the art. Alternatively, a secondary antibody specific for the antibodies of this invention is labeled with a known detectable label and used to detect the EphA2-specific antibodies in the above techniques. Thus, the antibodies of this invention
5 provide methods to detect metastatic transformation.

The present invention also employs antibodies specific for an extracellular epitope of EphA2 in therapeutic compositions and methods for use. When used for *in vivo* therapy, a pharmaceutical composition administered to a patient comprises EphA2-specific antibodies in therapeutically effective amounts in a
10 pharmaceutically acceptable carrier. In a preferred embodiment, the EphA2-specific antibodies have been "humanized." Humanized antibodies include "chimeric antibodies" made by splicing genes from a mouse (or other mammal) antibody of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity. Such techniques are known in the art. See, for
15 example, U.S. Patent No. 5,811,098. In addition to antibodies, natural or artificial ligands, peptides, anti-sense, ATP analogies, or other small molecules capable of specifically targeting EphA2 may be employed.

An example of another way to target EphA2 is the use of ephrins to activate or inhibit EphA2. For example, EphrinA1-F_c, the extracellular domain of
20 ephrinA1 linked to immunoglobulin heavy chain, increases the phosphotyrosine content of EphA2. EphrinA1-F_c reverses the malignant behavior of EphA2 transformed cells. Thus, another preferred embodiment of this invention is a pharmaceutical composition comprising an ephrin or a hybrid form of ephrin administered in a therapeutic amount.

25 Therapeutic amounts are amounts which eliminate or reduce the patient's tumor burden, or which prevent or reduce the proliferation of metastatic cells. The dosage will depend on many parameters, including the nature of the tumor, patient history, patient condition, the possible co-use of other oncolytic agents, and methods of administration. Methods of administration include injection (e.g.,
30 parenteral, subcutaneous, intravenous, intraperitoneal, etc.) for which the antibodies are provided in a nontoxic pharmaceutically acceptable carrier such as water, saline, Ringer's solution, dextrose solution, 5% human serum albumin, fixed oils, ethyl

oleate, or liposomes. Typical dosages may range from about 0.01 to about 20 mg/kg, and more particularly from about 0.1 to about 10 mg/kg. Other methods of administration include oral and transdermal. Acceptable carriers for oral ingestion in accordance with the present invention can be formulated using art-recognized techniques into pharmaceutically acceptable liquid carriers or in combination with pharmaceutically acceptable solid carriers in the form of tablets, capsules, caplets, or gel-seals. Other effective methods of administration and dosages may be determined by routine experimentation and are within the scope of this invention.

Therapeutic methods employing EphA2-specific antibodies may be combined with chemotherapy, surgery, and radiation therapy, depending on type of the tumor, patient condition, other health issues, and a variety of factors. The methods may also include immunoconjugates for targeted immunotoxin-mediated therapy, wherein antibodies of this invention are covalently or non-covalently conjugated to various cytotoxic agents, further enhancing toxicity to targeted cells. See, for example, U.S. Patent No. 5,872,223. Such agents, including various bacterial toxins (e.g., *Pseudomonas* exotoxin), ricin A-chain, daunorubicin, methotrexate, and ribosome inhibitors (e.g., trichosantin). Also, the antibodies of this invention may be labeled with alpha, beta, or Auger electron emitters, resulting in immunoconjugates for targeted radiotherapy.

Thus, EphA2-specific antibodies may be used in a variety of methods and compositions for detecting and treating metastatic disease.

EXAMPLE 1

Characterization of EphA2 Expression in Metastatic Cells

Following the RIMMS strategy using tyrosine phosphorylated proteins from Ras-transformed human epithelial cells, hybridomas were screened, and an antibody specific for EphA2 has been isolated. This antibody, was used to assess the levels of EphA2 expression in nontransformed prostatic epithelial cells and prostatic tumor cells. Low levels of EphA2 expression were found in non-transformed prostatic epithelial cells, but this EphA2 expression was enriched within sites of cell-cell contact and interacted with cell-bound ligand. Compared to non-transformed cells, two features distinguish EphA2 in metastatic prostate cancer cells: 1) EphA2 is

overexpressed; 2) EphA2 is diffusely distributed and does not appear to interact with ligand. To confirm these data, western blots were performed using the EphA2 specific antibodies. EphA2 overexpression in human prostate cancer cells (LNCAP, DU145, PC3) directly correlates with their invasiveness *in vitro* and *in vivo*. Of the
5 three lines tested, LNCAP is the least aggressive, DU145 is more aggressive, and PC3 is the most aggressive. As seen in Fig. 2, DU145 cells exhibit higher levels of EphA2 expression than LNCAP, and PC3 cells exhibit even higher levels of EphA2 expression. Similarly, as shown in Figs. 2B and 2C, EphA2 expression is elevated in variants of human prostatic epithelial cells transformed by oncogenic K-Ras or X-
10 irradiation. The three lanes in Fig. 2B show "normal" MCL prostatic epithelial cells, and K-Ras and X-ray transformed cell lines derived therefrom. Similarly, the three lanes of Fig. 2C show "normal" 267B1 prostatic epithelial cells, and K-Ras and X-ray transformed cell lines derived therefrom. As seen in Figs. 2B and 2C, the transformed cells all exhibited elevated EphA2 levels. Fig. 3 shows similar western blots, except
15 using prostate cancer cell lines from dogs. As shown in Fig. 3, consistent with the results from human cells, EphA2 is overexpressed in metastatic prostatic carcinoma cells derived from dogs with spontaneous prostate cancer.

The metastatic prostate cell lines can be subdivided into three categories: 1) cells derived from primary prostate tumors; 2) cells derived from
20 metastases that are poorly metastatic *in vivo*; 3) cells derived from metastases that are highly metastatic *in vivo*. The western blots using EphA2-specific antibodies have revealed that EphA2 expression is elevated in all cells derived from metastases, with highest EphA2 expression in cells that retain metastatic potential *in vivo* (as assessed using athymic mouse models). Interestingly, B2D6 studies have shown that EphA2 is
25 overexpressed in cells from prostate cancer metastases compared to lines established from the primary tumor of same patient. Taken together, these results all reveal EphA2 overexpression in metastatic prostate tumor cells.

Similar EphA2 expression patterns have been found with breast cancer cells. In normal mammary epithelial cells, EphA2 is enriched within the cell-cell
30 junctions. By contrast, non-metastatic breast cancer cells do not express EphA2, while metastatic breast cancer cells overexpress EphA2. In metastatic breast cancer

cells, EphA2 is redistributed into the membrane ruffles and, thus, is available for antibody binding.

EXAMPLE 2

5 *In Vitro* Targeting of Metastatic Cells

EphA2 overexpression renders metastatic cells susceptible to antibody-mediated selected killing with the present antibodies specific for an extracellular epitope of EphA2. While normal cells express EphA2, it is believed that ligand binding or clustering within sites of cell-cell contact occludes extracellular epitopes in
10 normal cells and renders them inaccessible to antibodies specific for an extracellular epitope of EphA2. The tumor selectivity of the antibodies of the present invention is believed to rival or exceed that of Herceptin® for targeting metastatic cancer.

EphA2 overexpression provides a basis for targeting metastatic cancer cells with EphA2-specific antibodies. Antibodies specific for an extracellular epitope
15 of EphA2, such as those produced by hybridoma B2D6, may be used to alter selectively (versus normal cells) the proliferative or invasive behaviors of metastatic cancer cells. In both metastasis-derived and laboratory-induced transformation, EphA2 overexpression correlates with invasiveness, whereas non-invasive cells have lower levels of EphA2 expression.

20 To measure the effect of B2D6 on cell growth, cells are incubated with purified B2D6, and cell proliferation is measured by counting cells microscopically (using a hemacytometer) and by measuring DNA synthesis. For example, normal growth media is supplemented with B2D6 and BrdU, and BrdU incorporation is measured over the following four hours. To measure the effects of B2D6 over longer
25 times, samples are counted at 24 hour intervals, with BrdU added to the culture media for the final four hours of incubation. As a third measure of cell growth, the effect of B2D6 on the growth of metastatic cells in soft agar is determined. Soft agar plating assays are used, wherein 2×10^4 metastatic cells are plated atop agar, in the presence or absence of B2D6 (0-10 nM), and colony growth is evaluated at three-day intervals
30 thereafter.

It is believed that B2D6 decreases the growth of metastatic cells. Preliminary results reveal that B2D6 aggregates EphA2 and blocks about 50% of

growth of metastatic breast cancer cells (which also overexpress EphA2) over the first four hours of incubation. Although EphA2 is not tyrosine phosphorylated in metastatic breast cancer cells, tyrosine phosphorylation is restored these B2D6 treated cells. Thus, B2D6 is believed to restore normal EphA2 function.

5 Additional studies with prostate cancer cells are being performed to determine if longer incubations with B2D6 further inhibit metastatic cell growth. Non-transformed epithelial cells express some EphA2, albeit much less than metastatic cells, and some toxicity to non-transformed cells is possible. The minimum effective and maximum non-toxic dosage levels of antibodies in accordance
10 with one aspect of this invention can be identified by routine experimentation, but preferably, typical doses will range from about 0.1 to about 20 mg/kg of patient body weight. The preferred dose will depend on many parameters, including the nature of the tumor, patient history, patient condition, the possible co-use of other oncolytic agents, and methods of administration. Antibody levels that best discriminate between
15 normal and metastatic cells will be used in treatment of metastatic tumors overexpressing EphA2 proteins.

EXAMPLE 3

In Vitro Antibody Mediated Cytotoxicity

20 Preliminary results demonstrate that EphA2 antibodies impede metastatic cell growth. To measure antibody-directed cytotoxicity, preferably, non-radioactive versions of ^{51}Cr -release assays using target cells (normal or metastatic prostatic epithelial cells) labeled with europium chloride (EuCl_3) and diethylenetriamine-pentaacetic acid (DTPA) are performed. After washing away
25 unincorporated Eu^{3+} , naphthoyltrifluoroacetone (NTA) and trioctylphosphine oxide are incubated with the cytolytic agents and assay supernatants. The luminescence of resultant ternary complex (Eu^{3+} /NTA/trioctylphosphine oxide) is measured using a fluorescence microplate reader. The sensitivity of this Eu^{3+} -release assay for complement-mediated cytotoxicity has been reported to be fivefold better than ^{51}Cr -
30 release assays. To determine specific lysis, parallel samples are hyponically lysed by adding distilled water. Untreated samples and isotype matched antibodies serve as negative controls.

To model complement-mediated death *in vitro*, cells are labeled with B2D6 and exposed to sera that has not been heat-inactivated. For antibody dependent cellular cytotoxicity ("ADCC"), both control and B2D6-treated prostate cells are incubated with peripheral blood mononuclear cells (PBMC; 10:1 E/T ratio).

5 It is believed that complement and ADCC both promote specific killing of the metastases. Antibody dose can be varied in order to establish LD₅₀ measurements for metastatic and normal cells. Antibody concentrations that maximize specific killing of metastatic cells (PC3, 267-Ras, 267-X) while minimizing the death to non-transformed prostatic epithelia (267, MLC) are determined by routine
10 experimentation. Also, ADCC can be combined with complement to further enhance tumor killing by treating the samples with tumor necrosis factor (TNF). There is evidence that TNF potentiates the killing of tumor cells by EGFR and Her2-specific antibodies. It is expected that humanizing the antibodies of this invention would provide for better complement or ADCC results.

15 It is believed that B2D6 kills tumor cells through complement cascade or ADCC. However, covalent or non-covalent conjugation of the present antibodies to art-recognized cytotoxic agents can further enhance toxicity to targeted cells. Examples of toxins appropriate for immunoconjugation include *Pseudomonas* exotoxin or ricin A-chain.

20

EXAMPLE 4

In Vivo targeting of Metastatic Cells

The present EphA2 antibodies, particularly those produced by hybridoma B2D6 are effective in blocking the growth and invasiveness of prostate
25 cancer cells *in vivo*. The efficacy of B2D6 in blocking the growth of primary prostate tumors using subcutaneous implantation of PC3 tumor cells in mice is determined by use of subcutaneous models. The primary advantages of subcutaneous models are the ease of implantation and subsequent monitoring of tumor size. 5x10⁵ PC3 cells are inoculated subcutaneously into the right craniolateral thorax (axilla) using aseptic
30 technique. Tumors are measured every 3-4 days using vernier calipers until they reach a volume of 0.2-0.3 cm³. At that time, the mice are divided into four groups (8-10 animals each): Group 1 (vehicle control), Groups 2-4 are treated with 0.1, 1.0, or

10 mg/kg B2D6, administered intraperitoneally, twice a week. The mice are then monitored every 3 days to measure tumor volume (with vernier calipers), body weight, and life span. After no greater than 60 days past implantation, the animals are sacrificed and postmortem evaluations of tumorigenesis, including measurement and weight of implanted tumors and proximal lymph nodes, macroscopic evaluation of soft tissues for tumors (lymph nodes and lung), and formalin fixation of the primary tumor and tissues, are performed. The tissues are evaluated by immunohistochemistry using D7 (another EphA2 specific antibody that is amenable to immunohistochemistry) to determine the level of EphA2 expression in the tumors. In particular, tumor cells that escape B2D6 treatment are studied to determine whether they have low levels of EphA2 expression. Also, EphA2 expression in the individual animals is correlated with tumor invasiveness.

As target-negative controls for specificity, parallel studies are performed using DU145 cells, which express very low levels of EphA2. Whereas the growth of PC3 tumors are believed to be sensitive to B2D6, tumors caused by DU145 are believed be insensitive. Statistical significance of B2D6 inhibition of tumorigenicity, overall metastatic frequency, and frequency of distant metastases is tested using computerized statistical packages (PC/SAS Ver. 6.04), with differences considered significant if $p < 0.05$.

While subcutaneous implantation is a popular and valuable method for modeling tumor cell growth, differences between the microenvironment of the skin and prostate can cause rather dramatic differences in cell behavior. For example, PC3 cells rarely metastasize when implanted subcutaneously whereas intraprostatic implantation (orthotopic) facilitates metastasis. Thus, PC3 cells are implanted in the prostate by exposing the prostate via laparotomy and inoculating tumor cells into prostate gland using a surgical microscope. After seven days, the mice begin receiving treatment with B2D6, as described above, and the animals are sacrificed no later than 60 days after implantation (or if the animals become moribund). The tumors are palpated at 3-5 day intervals, at which time data on tumor size, animal weight, and survival are collected. Post-mortem evaluations are also performed as described above, with emphasis upon the effect of B2D6 upon metastatic potential (to

lungs and regional lymph nodes). B2D6 is believed to block the primary tumor and metastatic potential of PC3 cells in a dose-dependent manner.

To minimize identification of strain or clonal-specific effects, identical analyses using other model systems can be employed. For example, the effects of B2D6 on the growth or metastasis of tumors caused by implantation of K-Ras or X-ray transformed 267B1 can be compared to the effects on MLC human prostatic epithelial cell lines.

EXAMPLE 5

10 **Development of "Second Generation" EphA2-based Antibody Therapeutics**

While EphA2 overexpression in metastatic prostate cancer cells provides a degree of selectivity comparable to Herceptin® in breast cancer, unique properties of EphA2 are believed to allow for even more selective targeting. In particular, EphA2 at the surface of non-transformed epithelia is tightly packed into cell-cell contacts whereas EphA2 on metastatic cells is diffusely distributed. It is thus likely that some epitopes on EphA2 are accessible in metastases but protected by ligand in normal prostatic epithelia. The EphA2 antibodies that provide the optimal discrimination between normal and metastatic prostatic epithelia are selected.

The panel of antibodies generated previously are screened for epitopes on EphA2 that are found at the cell surface. Using flow cytometry, EphA2 expression in a variety of "normal" (e.g., 267B or MLC cells) and metastatic cells (PC3 cells) are compared. For example, confluent monolayers of normal and metastatic cells are labeled with B2D6. To insure that antibodies are selected that are specific for epitopes inaccessible in normal cells but accessible in metastatic cells, antibodies are selected whose binding decreases in normal cells with increasing cell density but whose binding remains constant in metastatic cells, as shown in Fig. 4. After labeling with fluorescein-secondary antibodies, EphA2 expression is evaluated by using flow cytometry. The antibodies that best distinguish between normal and metastatic cells are selected. Specificity for EphA2 is confirmed via western blotting and immunoprecipitation studies. Antibodies exhibiting the best selectivity are then humanized using art recognized techniques.

EXAMPLE 6**Altered EphA2 Expression Through Transfection**

To assess the consequences of EphA2 overexpression, MCF-10A cells were transfected with human EphA2 cDNA (EphA2) or a vector control (vector).

- 5 After establishing cultures of MCF-10A cells with stable overexpression of EphA2, microscopic evaluation revealed differences in the cell morphology as compared to vector-transfected control cells (not shown). Non-transformed MCF-10A cells display an epithelial morphology and interact with one-another, even at low cell density. In contrast, EphA2-overexpressing MCF-10A cells (MCF^{EphA2}) adopt a
- 10 fibroblast-like morphology and do not form cell-cell contacts, even at high cell density. To confirm that the mesenchymal morphology does not represent clonal variation, a separate sample of MCF-10A cells transfected with EphA2 cDNAs yielded identical results.

- Cell-ECM adhesions were evaluated by incubating cells on ECM at
- 15 37°C for 30 minutes before vigorous washing to remove weakly adherent cells. These assays revealed a 24-fold increase in ECM attachments in MCF^{EphA2} cells relative to vector-transfected controls ($P < 4 \times 10^{-4}$). Cell-cell adhesions were assayed by incubating cells in suspension and counting the average size of cell colonies. Whereas vector-transfected MCF-10A cells interact with one-another in colonies with
- 20 an average size of 4.1 cells, the average colony size of MCF^{EphA2} cells is reduced to 1.3 cells ($P < 3 \times 10^{-5}$).

- Since stable cell-cell contacts cause EphA2 to become enriched within sites of cell-cell contact, EphA2 subcellular localization was assessed by immunostaining with specific antibodies. The EphA2 on non-transformed MCF-10A
- 25 cells was restricted to a narrow line where adjacent cells came into direct contact, with little staining of membrane that was not in contact with neighboring cells. In contrast, the pattern of EphA2 staining on MCF^{EphA2} cells was diffuse, with little staining of cell-cell contacts.

- The lack of EphA2 within cell-cell contacts in MCF^{EphA2} cells was
- 30 intriguing since EphA2 is stimulated by ligands that are anchored to the cell membrane. To measure EphA2 stimulation, the phosphotyrosine content of immunoprecipitated EphA2 was measured by Western blot analysis with

phosphotyrosine specific antibodies. Whereas the EphA2 in vector-transfected MCF-10 cells was tyrosine phosphorylated, EphA2 was not tyrosine phosphorylated in MCF^{EphA2} cells. The decreased phosphotyrosine content was confirmed using multiple EphA2 antibodies for immunoprecipitation (D7, B2D6) and different
5 phosphotyrosine-specific antibodies (4G10, PY20) for Western blot analyses.

EXAMPLE 7

Malignancy and Metastasis Through EphA2 Transfection

Malignant transformation was studied *in vitro*, and MCF^{EphA2} cells
10 were found to colonize soft agar. Whereas vector-transfected MCF-10A cells formed 0.3 colonies per high-power field, while MCF^{EphA2} cells displayed increased colony growth in soft agar, with an average of 3.0 colonies per high-power field ($P < 3 \times 10^{-7}$). Vector and EphA2 overexpressing MCF-10A cells were allowed to interact with Matrigel (Collaborative, Bedford, MA). Non-transformed MCF-10A cells rapidly
15 organized into spherical colonies when cultured on Matrigel whereas MCF^{EphA2} cells adopted a stellate organization that was indistinguishable from the behavior of metastatic cells (e.g., MDA-MB-231, MDA-MB,435).

Since *in vitro* analyses of transformation do not always predict tumorigenic potential *in vivo*, control or EphA2-overexpressing MCF-10A cells were
20 implanted into athymic (*nu/nu*) mice. Subcutaneous injection of MCF^{EphA2} cells caused the formation of palpable tumors within four days in 19 of 19 mice. The median volume of resulting tumors related to the number of implanted cells and reached an average of 300 mm³ (for samples injected with 5×10^6 cells) within 10 days (Table I). Necropsy revealed that the tumors were firmly attached to the underlying
25 axillary muscle and surrounded by fibrous tissue. Histologically, the neoplastic cells were invasive and associated with fibrous connective tissue. These neoplastic cells exhibited moderate cytoplasmic and nuclear pleiomorphism and formed dysplastic tubular and secreting structures. In control experiments, cells transfected with vector DNA failed to grow in athymic mice (0 of 13; Table I) and necropsy failed to identify
30 any growth or invasion of these cells.

Since the highest levels of EphA2 were consistently found in breast cancer cells that are metastatic *in vivo*, 1×10^6 control or MCF^{EphA2} cells were injected

into the tail vein of athymic mice. Within seven days, necropsy revealed lung micrometastases within large vessels in 2 of 4 mice injected with MCF^{EphA2} cells (Table I). The metastases were generally found to occlude large blood vessels but did not breach the vessel wall. Immunohistochemical staining with cytokeratin antibodies confirmed the epithelial nature of the thrombus and a lack of anti-thrombin staining revealed that the thrombus did not represent an abnormal or atypical outgrowth of endothelial cells. No lung colonization was observed in mice that had been injected with control MCF-10A cells (Table I).

Table I. Tumorigenic and Metastatic Potential of EphA2-Transformed MCF-10A Cells

Cell	Site of Inoculation	# of Cells Injected	Incidence of Tumorigenicity	Tumor Volume (mm ³)
Ctrl	Subcutaneous	1 x 10 ⁶	0/9	N/A
<i>EphA2</i>		1 x 10 ⁶	9/9	66 ± 20
Ctrl	Subcutaneous	5 x 10 ⁶	0/4	N/A
<i>EphA2</i>		5 x 10 ⁶	10/10	293 ± 70
Ctrl	Tail Vein	1 x 10 ⁶	0/4	
<i>EphA2</i>		1 x 10 ⁶	2/4	

EXAMPLE 8

Metastatic Targeting Using EphA2 Agonists

To test if EphA2 could be stimulated by an agonist, MCF^{EphA2} cells were suspended in soft agar in the presence or absence of 0.5 mg/mL EphrinA1-F_c. EphrinA1-F_c increased the phosphotyrosine content of EphA2, and EphrinA1-F_c-treated reduced colony formation in soft agar by 49% relative to vehicle-treated controls ($P < 5 \times 10^{-6}$). To test if EphA2 stimulation could alter cell behavior on Matrigel, the MCF^{EphA2} cells were treated with 0.5 mg/mL EphrinA1-F_c, which

restored a spherical phenotype that was comparable to non-transformed MCF-10A cells. Thus, EphA2 stimulation reverses the effects of EphA2 overexpression. EphrinA1-F_c. Despite its inability to interact with its endogenous ligands, the EphA2 in MCF^{EphA2} cells responded to exogenous stimuli.

- 5 Although the invention has been described in detail with reference to preferred embodiments, variations and modifications exist within the scope and spirit of the invention as described and defined in the following claims.

CLAIMS

1. A pharmaceutical composition for treatment of a mammalian metastatic tumor including a population of cells which overexpress tyrosine kinase
5 EphA2 having a peptide sequence defining at least one extracellular epitope of EphA2, said composition comprising a compound that specifically interacts with the extracellular epitope of EphA2 in an amount effective to reduce metastatic proliferation of said tumor, and a pharmaceutically acceptable carrier therefor.
2. The composition of claim 1 wherein the compound is an
10 antibody.
3. The composition of claim 2 wherein the antibody is a monoclonal antibody.
4. The composition of claim 3 wherein the antibody is the monoclonal antibody B2D6.
- 15 5. The composition of claim 2 wherein the antibody is conjugated to a cytotoxic agent.
6. The use of a compound for the manufacture of a medicament for the treatment of a metastatic tumor comprising a population of cells that overexpress EphA2, said compound having specificity for EphA2.
- 20 7. The use of claim 6 wherein the compound is an antibody having specificity for an extracellular epitope of EphA2.
8. The use of claim 7 wherein the antibody is a monoclonal antibody.
9. The use of claim 8 wherein the antibody is produced from a
25 hybridoma cell line identified as B2D6.
10. The use of claim 7 wherein the antibody selectively binds to metastatic cells.
11. The use of claim 10 wherein the antibody reduces proliferation of metastatic tumor cells.
- 30 12. The use of claim 6 wherein the population of cells forms at least a portion of a cancer tumor selected from the group consisting of breast, prostate, lung, and colon cancer tumors.

13. The use of claim 6 wherein the compound is an ephrin that affects phosphorylation of EphA2.

14. A method for detecting the presence of metastatic cells in a cell population comprising

5 providing an antibody specific to an extracellular epitope of EphA2,
binding a detectable label to the antibody,
providing a cell sample,
incubating the cell sample with the labeled antibody,
removing unbound antibody from the cell sample, and
10 detecting the presence of the label.

15. A method of producing an antibody which inhibits the proliferation of metastatic tumor cells comprising

injecting tyrosine phosphorylated proteins into lymph nodes of a
mammal,
15 harvesting lymph node cells from the mammal,
fusing the lymph node cells with myeloma cells to form hybridomas,
selecting at least one hybridoma producing an antibody which
specifically binds to an extracellular epitope of EphA2, and
isolating said antibody.

20 16. The method of claim 15 wherein the myeloma cells are Bcl-overexpressing.

17. The method of claim 15 wherein the tyrosine phosphorylated proteins comprise EphA2.

18. The antibody produced by the method of claim 15.

25 19. A pharmaceutical composition for treatment of a mammalian metastatic tumor, said composition comprising a compound that interferes with EphA2 function in an amount effective to reduce metastatic proliferation of said tumor, and a pharmaceutically acceptable carrier therefor.

30 20. The pharmaceutical composition of claim 19 wherein the compound is an agonist.

21. The pharmaceutical composition of claim 20 wherein the compound comprises a peptide sequence defining an extracellular domain of

EphrinA1.

22. The pharmaceutical composition of claim 21 wherein the peptide sequence is linked to a second peptide sequence defining immunoglobulin heavy chain.



- Inject Lymph Nodes Every Other Day for 10 Days
- Harvest Lymph Node Cells
- Fuse with Myeloma Overexpressing Bcl-2
- Screen by Immunofluorescence Microscopy
- Select for Antigen with Sites of Cell Adhesion:
 - Cell-Cell or
 - Cell-ECM/Membrane Ruffles

FIG. 1




FIG. 2A




FIG. 2B



FIG. 2C




FIG. 3




FIG. 4

Applicant's or agent's file reference number 290.01 J201	International application No. PCT/US00/22670
---	---

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13*bis*)

A. The indications made below relate to the microorganism referred to in the description on page <u>6</u> , line <u>7</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas VA 20110-2209 United States of America	
Date of deposit 8 December 2000 (08.12.00)	Accession Number PTA 2755
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Murine hybridoma D7	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

Applicant's or agent's file reference number 290.0. 0201	International application No. PCT/US00/22670
---	---

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>6</u> , line <u>8</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas VA 20110-2209 United States of America	
Date of deposit 8 December 2000 (08.12.00)	Accession Number PTA 2754
C. ADDITIONAL INDICATIONS (leave blank if not applicable)	
This information is continued on an additional sheet <input type="checkbox"/>	
Murine hybridoma B2D6	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
(Blank space for designated states)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
(Blank space for separate furnishing of indications)	

For receiving Office use only	
<input type="checkbox"/> This sheet was received with the international application	
Authorized officer	

For International Bureau use only	
<input type="checkbox"/> This sheet was received by the International Bureau on:	
Authorized officer	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22670

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/00, 39/395, 49/00; C07K 2/00, 5/00, 16/28

US CL : 424/9.2, 138.1, 141.1; 514/1 +; 530/350, 388.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/9.2, 138.1, 141.1; 514/1 +; 530/350, 388.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KINCH, M.S., et al, Identification of tyrosine phosphorylated adhesion proteins in human cancer cells, Hybridoma, 1998, Vol. 17, pages 227-235, especially Table 3, page 233.	1-12, 14-18
X,P	ZANTEK, N.C., et al, E-cadherin regulates the function of the EphA2 receptor tyrosine kinase, Cell Growth & Differentiation, September 1999, Vol. 10, pages 629-638, especially abstract and page 635, column 1.	1, 6, 12, 19-20
Y,P		1-12, 14-20
A	NAKAMOTO, M., Eph receptors and ephrins, International Journal of Biochemistry & Cell Biology, 1999, Vol. 32, pages 7-12.	1-22
A	HUAL, J., et al, Investigation of a possible receptor function of ephrinA ligands, European Journal of Neuroscience, 2000, Vol. 12, page 179.	1-14, 18-22
Y,P	WALKER-DANIELS, J., et al, Overexpression of the EphA2 tyrosine kinase in prostate cancer, The Prostate, 1999, Vol. 41, pages 275-280, especially abstract and page 276, column 1.	1-12, 14-18
Y	LINDBERG, R.A., et al, cDNA cloning and characterization of eck, an epithelial cell receptor protein-tyrosine kinase in the eph/elk family of protein kinases, Molecular and Cellular Biology, December 1990, Vol. 10, pages 6316-63254, especially abstract and page 6317, column 2.	1-12, 14-18

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

20 November 2000 (20.11.2000)

Date of mailing of the international search report

16 JAN 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Anthony C. Caputa, Ph.D.

Telephone No. (703) 308-0196

DELLA MAE COLLINS
PARALEGAL SPECIALIST
TECHNOLOGY CENTER 1600

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22670

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

- The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22670

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

- Species 1. An antibody specifically reactive to EphA2.
- Species 2. An ephrin capable of specific interaction with EphA2.
- Species 3. A peptide agonist of EphA2.

The claims are deemed to correspond to the species listed above in the following manner:

- Species 1. Claims 2-5, 7-11, 14-18
- Species 2. Claim 13
- Species 3. Claims 20-22

The following claim(s) are generic: claim 1, 6, 12, and 19.

The species listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:

The products of the above species represent separate and distinct molecules with different structures and functions such that one species could not be interchanged with the other. As such, each species would require different searches and the consideration of different patentability issues.

Continuation of B. FIELDS SEARCHED Item 3: STN/Medicine, Bioscience, Pharmacology, Toxicology
search terms: ephrin, EphA2, antibod?, antiser?, immuno?, Western, cancer, tyrosine kinase, metasta?, tumor, malignan?, monoclonal, polyclonal, therap?, treat?, inhibit?, hybridoma, B2D6, agonist, ligand, receptor, substrate, inhibitor, antagonist

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.